

# Higher order numerical methods for convective problems

Václav Kučera

Department of Numerical Mathematics  
Charles University in Prague

- 1 Standard FEM error estimates
  - Method of lines
  - Implicit scheme

- 1 Standard FEM error estimates
  - Method of lines
  - Implicit scheme

## Scalar nonlinear convection

$$\begin{aligned}
 \text{a)} \quad & \frac{\partial u}{\partial t} + \operatorname{div} \mathbf{f}(u) = g \\
 \text{b)} \quad & u|_{\Gamma_D \times (0, T)} = 0, \\
 \text{c)} \quad & \frac{\partial u}{\partial \mathbf{n}}|_{\Gamma_N \times (0, T)} = u_N, \\
 \text{d)} \quad & u(x, 0) = u^0(x), \quad x \in \Omega.
 \end{aligned}$$

- $\mathbf{f} \in [C_b^2(\mathbb{R})]^d$ ,
- $\mathbf{f}'(u) \cdot \mathbf{n} \geq 0$  on  $\Gamma_N$
- We assume  $u$  is sufficiently regular:

$$u, u_t \in L^2(0, T; H^{p+1}(\Omega))$$

- $p > \begin{cases} (d+1)/2, & \mathbf{f} \in [C_b^2(\mathbb{R})]^d, \\ (d-1)/2, & \mathbf{f} \in [C_b^3(\mathbb{R})]^d, \Gamma_N = \emptyset. \end{cases}$

## Scalar nonlinear convection

$$\text{a) } \frac{\partial u}{\partial t} + \operatorname{div} \mathbf{f}(u) = g$$

$$\text{b) } u|_{\Gamma_D \times (0, T)} = 0,$$

$$\text{c) } \frac{\partial u}{\partial \mathbf{n}}|_{\Gamma_N \times (0, T)} = u_N,$$

$$\text{d) } u(x, 0) = u^0(x), \quad x \in \Omega.$$

- $\mathbf{f} \in [C_b^2(\mathbb{R})]^d$ ,
- $\mathbf{f}'(u) \cdot \mathbf{n} \geq 0$  on  $\Gamma_N$
- We assume  $u$  is sufficiently regular:

$$u, u_t \in L^2(0, T; H^{p+1}(\Omega))$$

- $p > \begin{cases} (d+1)/2, & \mathbf{f} \in [C_b^2(\mathbb{R})]^d, \\ (d-1)/2, & \mathbf{f} \in [C_b^3(\mathbb{R})]^d, \Gamma_N = \emptyset. \end{cases}$

## Scalar nonlinear convection

$$\text{a) } \frac{\partial u}{\partial t} + \operatorname{div} \mathbf{f}(u) = g$$

$$\text{b) } u|_{\Gamma_D \times (0, T)} = 0,$$

$$\text{c) } \frac{\partial u}{\partial \mathbf{n}}|_{\Gamma_N \times (0, T)} = u_N,$$

$$\text{d) } u(x, 0) = u^0(x), \quad x \in \Omega.$$

- $\mathbf{f} \in [C_b^2(\mathbb{R})]^d$ ,
- $\mathbf{f}'(u) \cdot \mathbf{n} \geq 0$  on  $\Gamma_N$
- We assume  $u$  is sufficiently regular:

$$u, u_t \in L^2(0, T; H^{p+1}(\Omega))$$

- $p > \begin{cases} (d+1)/2, & \mathbf{f} \in [C_b^2(\mathbb{R})]^d, \\ (d-1)/2, & \mathbf{f} \in [C_b^3(\mathbb{R})]^d, \Gamma_N = \emptyset. \end{cases}$

## Scalar nonlinear convection

$$\text{a) } \frac{\partial u}{\partial t} + \operatorname{div} \mathbf{f}(u) = g$$

$$\text{b) } u|_{\Gamma_D \times (0, T)} = 0,$$

$$\text{c) } \frac{\partial u}{\partial \mathbf{n}}|_{\Gamma_N \times (0, T)} = u_N,$$

$$\text{d) } u(x, 0) = u^0(x), \quad x \in \Omega.$$

- $\mathbf{f} \in [C_b^2(\mathbb{R})]^d$ ,
- $\mathbf{f}'(u) \cdot \mathbf{n} \geq 0$  on  $\Gamma_N$
- We assume  $u$  is sufficiently regular:

$$u, u_t \in L^2(0, T; H^{p+1}(\Omega))$$

- $p > \begin{cases} (d+1)/2, & \mathbf{f} \in [C_b^2(\mathbb{R})]^d, \\ (d-1)/2, & \mathbf{f} \in [C_b^3(\mathbb{R})]^d, \Gamma_N = \emptyset. \end{cases}$

## Scalar nonlinear convection

$$\text{a) } \frac{\partial u}{\partial t} + \operatorname{div} \mathbf{f}(u) = g$$

$$\text{b) } u|_{\Gamma_D \times (0, T)} = 0,$$

$$\text{c) } \frac{\partial u}{\partial \mathbf{n}}|_{\Gamma_N \times (0, T)} = u_N,$$

$$\text{d) } u(x, 0) = u^0(x), \quad x \in \Omega.$$

- $\mathbf{f} \in [C_b^2(\mathbb{R})]^d$ ,
- $\mathbf{f}'(u) \cdot \mathbf{n} \geq 0$  on  $\Gamma_N$
- We assume  $u$  is sufficiently regular:

$$u, u_t \in L^2(0, T; H^{p+1}(\Omega))$$

- $p > \begin{cases} (d+1)/2, & \mathbf{f} \in [C_b^2(\mathbb{R})]^d, \\ (d-1)/2, & \mathbf{f} \in [C_b^3(\mathbb{R})]^d, \Gamma_N = \emptyset. \end{cases}$



## Definition

Standard conforming  $p$ -order FEM solution of the convection-diffusion problem:

a)  $u_h \in C^1([0, T]; V_h),$

b)  $\left( \frac{\partial u_h(t)}{\partial t}, \varphi_h \right) + \mathbf{b}(u_h(t), \varphi_h) = \ell(\varphi_h)(t), \quad \forall \varphi_h \in V_h, \forall t \in (0, T),$

c)  $u_h(0) = u_h^0.$

Convective term

$$\mathbf{b}(u, v) = - \int_{\Omega} \mathbf{f}(u) \cdot \nabla v \, dx$$

## Definition

Standard conforming  $p$ -order FEM solution of the convection-diffusion problem:

a)  $u_h \in C^1([0, T]; V_h),$

b)  $\left( \frac{\partial u_h(t)}{\partial t}, \varphi_h \right) + b(u_h(t), \varphi_h) = \ell(\varphi_h)(t), \quad \forall \varphi_h \in V_h, \forall t \in (0, T),$

c)  $u_h(0) = u_h^0.$

Right-hand side term

$$\ell(\mathbf{v})(t) = \int_{\Omega} g(t) \mathbf{v} \, dx$$

# Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

# Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

# Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

## Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

$$\underbrace{\left( \frac{d\xi}{dt}, \xi \right)}_{\frac{1}{2} \frac{d}{dt} \|\xi(t)\|^2} = b(u_h, \xi) - b(u, \xi) + \left( \frac{d\eta}{dt}, \xi \right)$$

## Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

$$\left( \frac{d\xi}{dt}, \xi \right) = b(u_h, \xi) - b(u, \xi) + \underbrace{\left( \frac{d\eta}{dt}, \xi \right)}_{\leq O(h^{2p+2}) + \|\xi\|^2}$$

## Error estimates

- Let  $e_h = \eta + \xi$ , where  $\eta = \Pi_h u - u$ ,  $\xi = u_h - \Pi_h u \in V_h$ .
- $\Pi_h : L^2(\Omega) \rightarrow V_h$  is the  $L^2(\Omega)$ -projection
- $\eta = O(h^\mu)$  in various norms,  $\xi = ?$
- Subtract  $eq(u) - eq(u_h)$ , set  $\varphi_h := \xi$

$$\left( \frac{d\xi}{dt}, \xi \right) = b(u_h, \xi) - b(u, \xi) + \left( \frac{d\eta}{dt}, \xi \right)$$

$$\Downarrow$$

$$\frac{d}{dt} \|\xi(t)\|^2 \leq b(u_h, \xi) - b(u, \xi) + O(h^{2p+2}) + \|\xi\|^2$$



$$\frac{d}{dt} \|\xi(t)\|^2 \leq b(u_h, \xi) - b(u, \xi) + O(h^{2p+2}) + \|\xi\|^2$$

- For Gronwall we need only  $h^{2p+2}$ ,  $\|\xi\|^2$  on the RHS. Then

$$\max_{t \in [0, T]} \|\xi(t)\|^2 dt = O(h^{2p+2}).$$

- Naively

$$b(u_h, \xi) - b(u, \xi) = \int_{\Omega} (\mathbf{f}(u) - \mathbf{f}(u_h)) \cdot \nabla \xi \, dx \leq C \|e_h\| \|\xi\|_1 \leq \frac{C}{\varepsilon} \|e_h\|^2 + \frac{1}{2} \varepsilon \|\xi\|_1^2,$$

- If we estimate using the inverse inequality

$$b(u_h, \xi) - b(u, \xi) \leq C \|e_h\| \|\xi\|_1 \leq C \|e_h\| C_I h^{-1} \|\xi\|,$$

then we get  $O(\exp(\frac{C}{h}) h^{2p+2})$ .

$$\frac{d}{dt} \|\xi(t)\|^2 \leq b(u_h, \xi) - b(u, \xi) + O(h^{2p+2}) + \|\xi\|^2$$

- For Gronwall we need only  $h^{2p+2}$ ,  $\|\xi\|^2$  on the RHS. Then

$$\max_{t \in [0, T]} \|\xi(t)\|^2 dt = O(h^{2p+2}).$$

- Naively

$$b(u_h, \xi) - b(u, \xi) = \int_{\Omega} (\mathbf{f}(u) - \mathbf{f}(u_h)) \cdot \nabla \xi \, dx \leq C \|e_h\| \|\xi\|_1 \leq \frac{C}{\varepsilon} \|e_h\|^2 + \frac{1}{2} \varepsilon \|\xi\|_1^2,$$

- If we estimate using the inverse inequality

$$b(u_h, \xi) - b(u, \xi) \leq C \|e_h\| \|\xi\|_1 \leq C \|e_h\| C_I h^{-1} \|\xi\|,$$

then we get  $O(\exp(\frac{C}{h}) h^{2p+2})$ .

$$\frac{d}{dt} \|\xi(t)\|^2 \leq b(u_h, \xi) - b(u, \xi) + O(h^{2p+2}) + \|\xi\|^2$$

- For Gronwall we need only  $h^{2p+2}$ ,  $\|\xi\|^2$  on the RHS. Then

$$\max_{t \in [0, T]} \|\xi(t)\|^2 dt = O(h^{2p+2}).$$

- Naively

$$b(u_h, \xi) - b(u, \xi) = \int_{\Omega} (\mathbf{f}(u) - \mathbf{f}(u_h)) \cdot \nabla \xi \, dx \leq C \|e_h\| \|\xi\|_1 \leq \frac{C}{\varepsilon} \|e_h\|^2 + \frac{1}{2} \varepsilon \|\xi\|_1^2,$$

- If we estimate using the inverse inequality

$$b(u_h, \xi) - b(u, \xi) \leq C \|e_h\| \|\xi\|_1 \leq C \|e_h\| C_I h^{-1} \|\xi\|,$$

then we get  $O(\exp(\frac{c}{h}) h^{2p+2})$ .

The estimate of *Zhang, Shu (2004)*

## Lemma

$$b(u_h, \xi) - b(u, \xi) \leq C \left( 1 + \frac{\|e_h\|_\infty}{h} \right) (h^{2p+1} + \|\xi\|^2)$$

- If  $\mathbf{f} \in [C_b^3(\mathbb{R})]^d$ , then we get a factor of  $\frac{\|e_h\|_\infty^2}{h}$
- If  $\|e_h(t)\|_\infty = O(h)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

- For an explicit scheme, *Zhang, Shu (2004)* use induction:

$$\|e_h(t_n)\| = O(h^{p+1/2}) \Rightarrow \|e_h(t_{n+1})\|_\infty = O(h) \Rightarrow \|e_h(t_{n+1})\| = O(h^{p+1/2})$$

The estimate of *Zhang, Shu (2004)*

## Lemma

$$b(u_h, \xi) - b(u, \xi) \leq C \left( 1 + \frac{\|e_h\|_\infty}{h} \right) (h^{2p+1} + \|\xi\|^2)$$

- If  $\mathbf{f} \in [C_b^3(\mathbb{R})]^d$ , then we get a factor of  $\frac{\|e_h\|_\infty^2}{h}$
- If  $\|e_h(t)\|_\infty = O(h)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C (h^{2p+1} + \|\xi\|^2).$$

- For an explicit scheme, *Zhang, Shu (2004)* use induction:

$$\|e_h(t_n)\| = O(h^{p+1/2}) \Rightarrow \|e_h(t_{n+1})\|_\infty = O(h) \Rightarrow \|e_h(t_{n+1})\| = O(h^{p+1/2})$$

The estimate of *Zhang, Shu (2004)*

## Lemma

$$b(u_h, \xi) - b(u, \xi) \leq C \left( 1 + \frac{\|e_h\|_\infty}{h} \right) (h^{2p+1} + \|\xi\|^2)$$

- If  $\mathbf{f} \in [C_b^3(\mathbb{R})]^d$ , then we get a factor of  $\frac{\|e_h\|_\infty^2}{h}$
- If  $\|e_h(t)\|_\infty = O(h)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

- For an explicit scheme, *Zhang, Shu (2004)* use induction:

$$\|e_h(t_n)\| = O(h^{p+1/2}) \Rightarrow \|e_h(t_{n+1})\|_\infty = O(h) \Rightarrow \|e_h(t_{n+1})\| = O(h^{p+1/2})$$

The estimate of *Zhang, Shu (2004)*

## Lemma

$$b(u_h, \xi) - b(u, \xi) \leq C \left( 1 + \frac{\|e_h\|_\infty}{h} \right) (h^{2p+1} + \|\xi\|^2)$$

- If  $\mathbf{f} \in [C_b^3(\mathbb{R})]^d$ , then we get a factor of  $\frac{\|e_h\|_\infty^2}{h}$
- If  $\|e_h(t)\|_\infty = O(h)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

- For an explicit scheme, *Zhang, Shu (2004)* use induction:

$$\|e_h(t_n)\| = O(h^{p+1/2}) \Rightarrow \|e_h(t_{n+1})\|_\infty = O(h) \Rightarrow \|e_h(t_{n+1})\| = O(h^{p+1/2})$$

## Lemma

If  $\|e_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .

## Main theorem

Let  $p > (d+1)/2$ . Then

$$\|e_h\|_{L^\infty(L^2)} \leq C_T h^{p+1/2},$$

*Proof:*

- Nonlinear Gronwall-type lemma.



## Lemma

If  $\|e_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .

## Main theorem

Let  $p > (d+1)/2$ . Then

$$\|e_h\|_{L^\infty(L^2)} \leq C_T h^{p+1/2},$$

*Proof:*

- Nonlinear Gronwall-type lemma.
- Continuous mathematical induction (Y. R. Chao, 1919)

## Lemma

If  $\|e_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .

## Main theorem

Let  $p > (d+1)/2$ . Then

$$\|e_h\|_{L^\infty(L^2)} \leq C_T h^{p+1/2},$$

*Proof:*

- Nonlinear Gronwall-type lemma.
- Continuous mathematical induction (Y. R. Chao, 1919)

## Lemma

If  $\|e_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .


## Main theorem


Let  $p > (d+1)/2$ . Then


$$\|e_h\|_{L^\infty(L^2)} \leq C_T h^{p+1/2},$$


*Proof:*


- Nonlinear Gronwall-type lemma.
- Continuous mathematical induction (Y. R. Chao, 1919)
















- 1 Standard FEM error estimates
  - Method of lines
  - Implicit scheme

## Definition

Let  $0 = t_0 < t_1 < \dots < t_{N+1} = T$ ,  $\tau_n := t_{n+1} - t_n$

a)  $u_h^n \in V_h$ ,

b)  $\left( \frac{u_h^{n+1} - u_h^n}{\tau_n}, \varphi_h \right) + b(u_h^{n+1}, \varphi_h) = \ell(\varphi_h)(t_{n+1}),$

$$\forall \varphi_h \in V_h, \forall n = 0, \dots, N,$$

c)  $u_h^0 \approx u(0).$

# Standard approach

- $eq(u) - eq(u_h)$
- test by  $\xi$
- estimate  $b, \ell$
- use Gronwall's inequality.

## Theorem

*There does not exist a Gronwall type lemma which could prove the desired error estimate only from the error equation tested by  $\xi$  and estimates of individual terms contained therein.*

# Standard approach

- $eq(u) - eq(u_h)$
- test by  $\xi$
- estimate  $b, \ell$
- use Gronwall's inequality.

## Theorem

*There does not exist a Gronwall type lemma which could prove the desired error estimate only from the error equation tested by  $\xi$  and estimates of individual terms contained therein.*

# Standard approach

- $eq(u) - eq(u_h)$
- test by  $\xi$
- estimate  $b, \ell$
- use Gronwall's inequality.

## Theorem

*There does not exist a Gronwall type lemma which could prove the desired error estimate only from the error equation tested by  $\xi$  and estimates of individual terms contained therein.*

# Standard approach

- $eq(u) - eq(u_h)$
- test by  $\xi$
- estimate  $b, \ell$
- use Gronwall's inequality.

## Theorem

*There does not exist a Gronwall type lemma which could prove the desired error estimate only from the error equation tested by  $\xi$  and estimates of individual terms contained therein.*

# Standard approach

- $eq(u) - eq(u_h)$
- test by  $\xi$
- estimate  $b, \ell$
- use Gronwall's inequality.

## Theorem

*There does not exist a Gronwall type lemma which could prove the desired error estimate only from the error equation tested by  $\xi$  and estimates of individual terms contained therein.*



## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_h^{n+1} - u_h^n}{\tau_n}, \varphi_h \right) + b(u_h^{n+1}, \varphi_h) = \ell(\varphi_h)(t_{n+1}), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

Lemma (Existence, uniqueness and continuity)

Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .

Definition (Continuated discrete solution)

Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_\tau - U_h}{\tau}, \varphi_h \right) + b(u_\tau, \varphi_h) = l(\varphi_h)(t), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

Lemma (Existence, uniqueness and continuity)

*Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .*

Definition (Continuated discrete solution)

Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_\tau - U_h}{\tau}, \varphi_h \right) + b(u_\tau, \varphi_h) = l(\varphi_h)(t), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

Lemma (Existence, uniqueness and continuity)

*Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .*

Definition (Continuated discrete solution)

Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_\tau - U_h}{\tau}, \varphi_h \right) + b(u_\tau, \varphi_h) = l(\varphi_h)(t), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

Lemma (Existence, uniqueness and continuity)

*Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .*

Definition (Continuated discrete solution)

Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_\tau - U_h}{\tau}, \varphi_h \right) + b(u_\tau, \varphi_h) = l(\varphi_h)(t), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

## Lemma (Existence, uniqueness and continuity)

Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .

## Definition (Continuated discrete solution)

Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation

## Auxiliary problem

Given  $\tau \geq 0$  and  $U_h \in V_h$ , we seek  $u_\tau \in V_h$  such that

$$\left( \frac{u_\tau - U_h}{\tau}, \varphi_h \right) + b(u_\tau, \varphi_h) = l(\varphi_h)(t), \quad \forall \varphi_h \in V_h.$$

By setting  $U_h := u_h^n$ ,  $\tau := \tau_n$ , then  $u_\tau = u_h^{n+1}$ .

By setting  $U_h := u_h^n$ ,  $\tau = 0$ , then  $u_\tau = u_h^n$ .

## Lemma (Existence, uniqueness and continuity)


Let  $\tau = O(h)$ , then  $\exists! u_\tau \in V_h$  and  $\|u_\tau\|$  depends continuously on  $\tau$ .

## Definition (Continuated discrete solution)


Let  $\tilde{u}_h : [0, T] \rightarrow V_h$  be such that for  $t \in [t_n, t_{n+1}]$  we define  $\tilde{u}_h(t) := u_\tau$ , the solution of the auxiliary problem with  $\tau := t - t_n$  and  $U_h := u_h^n$ .

Furthermore, we define  $\tilde{e}_h := u - \tilde{u}_h$ .

## Continuation




## Continuation






## Continuation



# Continuation



## Remark

Estimates for  $\tilde{e}_h \implies$  Estimates for  $e_h^n$ ,  $n = 0, \dots, N + 1$ .

## Lemma

If  $\|\tilde{e}_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|\tilde{e}_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

## Main theorem

Let  $p > (d + 1)/2$  and  $\tau = O(h^{1+d/2})$ . Then

$$\|\tilde{e}_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

*Proof:* Continuous mathematical induction.

## Remark

Estimates for  $\tilde{e}_h \implies$  Estimates for  $e_h^n$ ,  $n = 0, \dots, N + 1$ .

## Lemma

If  $\|\tilde{e}_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|\tilde{e}_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

## Main theorem

Let  $p > (d + 1)/2$  and  $\tau = O(h^{1+d/2})$ . Then

$$\|\tilde{e}_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

*Proof:* Continuous mathematical induction.

## Remark

Estimates for  $\tilde{e}_h \implies$  Estimates for  $e_h^n$ ,  $n = 0, \dots, N + 1$ .

## Lemma

If  $\|\tilde{e}_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|\tilde{e}_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

## Main theorem

Let  $p > (d + 1)/2$  and  $\tau = O(h^{1+d/2})$ . Then

$$\|\tilde{e}_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

*Proof:* Continuous mathematical induction.

## Remark

Estimates for  $\tilde{e}_h \implies$  Estimates for  $e_h^n$ ,  $n = 0, \dots, N + 1$ .

## Lemma

If  $\|\tilde{e}_h(\vartheta)\|_\infty = O(h)$  for all  $\vartheta \in (0, t)$ , then

$$\|\tilde{e}_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

## Main theorem

Let  $p > (d + 1)/2$  and  $\tau = O(h^{1+d/2})$ . Then

$$\|\tilde{e}_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C(h^{p+1/2} + \tau),$$

*Proof:* Continuous mathematical induction.

# From *globally* to *locally* Lipschitz $\mathbf{f}$

- We assume only  $\mathbf{f} \in (C^2(\mathbb{R}))^d$ .
- Zhang & Shu modify  $\mathbf{f}$  far from  $\mathcal{R}(u)$  to obtain  $\mathbf{f} \in (C_b^2(\mathbb{R}))^d$ . This does not change  $u$ , but we get a completely new scheme.
- We prove error estimates directly for locally Lipschitz  $\mathbf{f}$ .

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

- We assume only  $\mathbf{f} \in (C^2(\mathbb{R}))^d$ .
- Zhang & Shu modify  $\mathbf{f}$  far from  $\mathcal{R}(u)$  to obtain  $\mathbf{f} \in (C_b^2(\mathbb{R}))^d$ . This does not change  $u$ , but we get a completely new scheme.
- We prove error estimates directly for locally Lipschitz  $\mathbf{f}$ .



# From *globally* to *locally* Lipschitz $\mathbf{f}$

- We assume only  $\mathbf{f} \in (C^2(\mathbb{R}))^d$ .
- Zhang & Shu modify  $\mathbf{f}$  far from  $\mathcal{R}(u)$  to obtain  $\mathbf{f} \in (C_b^2(\mathbb{R}))^d$ . This does not change  $u$ , but we get a completely new scheme.
- We prove error estimates directly for locally Lipschitz  $\mathbf{f}$ .

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

- Let  $h \in (0, h_0)$ ,  $t \in [0, T]$ . We define the *admissible set*  $\mathcal{U}_h^{ad}(t) := \{v \in \mathcal{S}_h; \|u(t) - v\| \leq h^{1+d/2}\}$ .
- Functions from  $\mathcal{U}_h^{ad}(t)$  have values in some fixed compact  $[-R; R]$ . Hence,  $\mathbf{f}$  is Lipschitz continuous on  $[-R; R]$ .

## Lemma

If  $u_h(t) \in \mathcal{U}_h^{ad}(t)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

- Let  $h \in (0, h_0)$ ,  $t \in [0, T]$ . We define the *admissible set*  $\mathcal{U}_h^{ad}(t) := \{v \in \mathcal{S}_h; \|u(t) - v\| \leq h^{1+d/2}\}$ .
- Functions from  $\mathcal{U}_h^{ad}(t)$  have values in some fixed compact  $[-R; R]$ . Hence,  $\mathbf{f}$  is Lipschitz continuous on  $[-R; R]$ .

## Lemma

If  $u_h(t) \in \mathcal{U}_h^{ad}(t)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

- Let  $h \in (0, h_0)$ ,  $t \in [0, T]$ . We define the *admissible set*  $\mathcal{U}_h^{ad}(t) := \{v \in \mathcal{S}_h; \|u(t) - v\| \leq h^{1+d/2}\}$ .
- Functions from  $\mathcal{U}_h^{ad}(t)$  have values in some fixed compact  $[-R; R]$ . Hence,  $\mathbf{f}$  is Lipschitz continuous on  $[-R; R]$ .

## Lemma

If  $u_h(t) \in \mathcal{U}_h^{ad}(t)$ , then

$$b_h(u_h, \xi) - b_h(u, \xi) \leq C(h^{2p+1} + \|\xi\|^2).$$

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

## Lemma

If  $u_h(\vartheta) \in \mathcal{U}_h^{ad}(\vartheta)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .

## Main theorem

Let  $p > 1 + d/2$ . Then

$$\|e_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C_T h^{p+1/2},$$

*Proof:* Continuous mathematical induction.

From *globally* to *locally* Lipschitz  $\mathbf{f}$ 

## Lemma

If  $u_h(\vartheta) \in \mathcal{U}_h^{ad}(\vartheta)$  for all  $\vartheta \in (0, t)$ , then

$$\|e_h\|_{L^\infty(0,t;L^2(\Omega))} \leq C_T h^{p+1/2},$$

where  $C_T$  is independent of  $h, t$ .

## Main theorem

Let  $p > 1 + d/2$ . Then

$$\|e_h\|_{L^\infty(0,T;L^2(\Omega))} \leq C_T h^{p+1/2},$$

*Proof:* Continuous mathematical induction.

# Conclusions and outlook

- Estimates for **nonlinear** convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?

# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?



# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?

# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?

# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for **locally** Lipschitz nonlinearities.
  - Possible (but technical) for DG.
  - Incompressible N-S convective term?

# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?

# Conclusions and outlook

- Estimates for nonlinear convection equations under high regularity assumptions.
- Analysis is valid for higher order elements:  $p > (d - 1)/2$ .
- Unnatural CFL condition  $\tau = O(h^{(1+d)/2})$  for implicit case.
- The situation would improve for higher order discretizations in time, e.g. BDF, space-time DG, ... CFL condition  $\tau = O(h^{(1+d)/2k})$  for a  $k$ -th order scheme in time.
- Estimates for locally Lipschitz nonlinearities.
- Possible (but technical) for DG.
- Incompressible N-S convective term?

Thank you for your attention.